

Integrilin (eptifibatide)

Indications Percutaneously Coronary Intervention (PCI) & Acute Coronary Syndrome
 Loading Dose 180 mcg/kg intravenous bolus
 Repeat Loading Dose For PCI only repeat loading dose 10 minutes after 1st loading dose

Maintenance Infusion 2 mcg/kg/min: Creatinine clearance greater than or equal to 50 ml/min or if creatinine clearance is unavailable a serum creatinine less than or equal to 2 mg/dl
 1 mcg/kg/min: Creatinine clearance less than 50 ml/min or if creatinine clearance is unavailable a Serum Creatinine greater than 2 mg/dl
 A creatinine clearance calculator is on the pharmacy web site.

Incompatible drugs Furosemide
 Compatible drugs atropine, dobutamine, heparin, lidocaine, meperidine, metoprolol, midazolam
 Compatible drugs morphine, nitroglycerin, potassium, verapamil

Contraindications bleeding diathesis, HTN (SBP > 200 DBP > 110), major surgery within 6 weeks, history of stroke within 30 days or any history of hemorrhagic stroke, dependency on renal dialysis, hypersensitivity to component of product

Weight	Loading Dose From 2 mg/ml Vial	Maintenance Infusion From 0.75 mg/ml Bottle	
		2 mcg/kg/min for Creatinine clearance greater than or equal to 50 ml/min or if creatinine clearance is unavailable a Serum Creatinine less than or equal to 2 mg/dl	1 mcg/kg/min for Creatinine clearance less than 50 ml/min or if creatinine clearance is unavailable a Serum Creatinine greater than 2 mg/dl
40	3.6	6.4	3.2
45	4.1	7.2	3.6
50	4.5	8	4
55	5.0	8.8	4.4
60	5.4	9.6	4.8
65	5.9	10.4	5.2
70	6.3	11.2	5.6
75	6.8	12	6
80	7.2	12.8	6.4
85	7.7	13.6	6.8
90	8.1	14.4	7.2
95	8.6	15.2	7.6
100	9.0	16	8
105	9.5	16.8	8.4
110	9.9	17.6	8.8
115	10.4	18.4	9.2
120	10.8	19.2	9.6
121	11.3	20	10
>121	11.3	20	10
kg	ml	ml/hr	