



BON SECOURS
RICHMOND HEALTH SYSTEM

- Memorial Regional Medical Center
- Richmond Community Hospital
- St. Francis Medical Center
- St. Mary's Hospital

Place patient label inside box (if no patient label, complete below)

Name: _____

DOB: _____

MR #: _____

**ARGATROBAN PROTOCOL
FOR HIT/ISOLATED HIT**

ALL ORDERS MUST INCLUDE DATE, TIME,
AND PHYSICIAN'S SIGNATURE.

THIS PATIENT IS ALLERGIC TO

NKDA

Date: _____ **Estimated Length of Stay:** _____

ORDER PATIENT STATUS: **Observation Indication** _____ (e.g., nausea, bleeding) **Outpatient in a bed**
(additional time needed to evaluate patient's condition) (Normal recovery requiring acute care bed)

Inpatient (meets medical necessity for admission) **Outpatient procedure**

Hospitalization certified necessary for the following reasons:

Plan of Care: See Progress Notes, H and P, and Physician Orders

D/C Plan: Home with Office Follow-up Home Health Care Rehabilitation Care Facility
 Extended Care Facility (e.g. Adult Home, Nursing Home, etc.) Other (Specify): _____

Physician Signature: _____

ORDERS (other than medications) CHECK ALL THAT APPLY	MEDICATION & I.V. FLUID ORDERS CHECK ALL THAT APPLY
<ol style="list-style-type: none"> 1. Patient must be in critical care or step down unit 2. Baseline PTT, PT/INR, Creatinine, CMP, HCT, Platelet Count 3. Platelet count every day 4. All PTTs are to be run stat 5. PTT 2 hr and 6 hr after initiation of Argatroban 6. PTT 4 hr after any dosage change until two consecutive PTTs are therapeutic (40-83 seconds) then change PTT to q day 7. If PTT is greater than 100 seconds stop argatroban, draw PTT every 2 hours until PTT is 40-83 seconds then restart Argatroban at new rate (see Rate Chart) 8. Daily PT/INR after warfarin started as noted below <ol style="list-style-type: none"> a. If Argatroban dose is greater than 2 mcg/kg/min, reduce Argatroban to 2 mcg/kg/min for 4 hours, draw PT/INR. If PT/INR greater than 4, on Warfarin and Argatroban, stop Argatroban, draw PT/INR four hours later, restart Argatroban immediately after PT/INR drawn at original rate. Use second PT/INR for Warfarin dosage adjustments. b. If Argatroban dose is less than or equal to 2 mcg/kg/min drawn PT/INR, if PT/INR greater than 4, on Warfarin and Argatroban, stop Argatroban, draw PT/INR four hours later, restart Argatroban immediately after PT/INR drawn at prior rate. Use second PT/INR for Warfarin dosage adjustments. 9. Guaiac all stools, gastrocuccult all emesis, visually check for hematuria or other bleeding. 10. Notify physician immediately for unexplained drop in blood pressure, development of hematoma, drop in HCT, flank pain, bright red urine, bruising or other signs of bleeding. 	<p>Discontinue all Heparin, Heparin flushes, Enoxaparin (Lovenox) and Dalteparin (Fragmin)</p> <p>Discontinue Warfarin (Coumadin)</p> <p>Phytonadione (Vitamin K) 10 mg subQ x 1 (if the patient has been on Warfarin)</p> <p>Discontinue all IM injections (physician to give order for alternate route of administration)</p> <p>Argatroban infusion:</p> <p><input type="checkbox"/> 2 mcg/kg/min if hepatic disease score per table is less than or equal to 6</p> <p><input type="checkbox"/> 1 mcg/kg/min if creatinine clearance is less than 60 ml/min and hepatic disease score is less than or equal to 6.</p> <p><input type="checkbox"/> 0.5 mcg/kg/min if hepatic disease score per table is greater than 6</p> <p>Adjust Argatroban rate per PTT using Argatroban rate adjustment chart</p> <p>Pharmacy to send initial infusion rate chart and rate adjustment chart to floor with Argatroban infusion</p> <p>Warfarin (Coumadin) _____ mg (range 2 - 5 mg) Qday once platelet count is greater than 150,000 per mcL Goal PT/INR while argatroban infusing at or below 2 mcg/kg/min is 4-5, off argatroban goal PT/INR is 2-3.</p> <p>Discontinue Argatroban when two consecutive daily INR are 2 to 3 (on Warfarin alone, see step 8 on left) and Warfarin therapy has overlapped Argatroban therapy for at least 5 days.</p>

MR-PHY-6-79 (02/07)

(SMH TAB # 2)

DO NOT USE ABBREVIATIONS:			
MS	U (unit)	IU (International Unit)	Q.D. QD q.d. qd (daily)
MISO4	Trailing zero (X.0mg)		Q.O.D. QOD q.o.d. qod (every other day)
MgSO4	Lack of leading zero (.Xmg)		

Physician Signature _____ Date / Time _____



BON SECOURS
RICHMOND HEALTH SYSTEM

- Memorial Regional Medical Center
- Richmond Community Hospital
- St. Francis Medical Center
- St. Mary's Hospital

Place patient label inside box (if no patient label, complete below)

Name: _____

DOB: _____

MR #: _____

**ARGATROBAN PROTOCOL
FOR HIT/ISOLATED HIT**

ALL ORDERS MUST INCLUDE DATE, TIME,
AND PHYSICIAN'S SIGNATURE.

THIS PATIENT IS ALLERGIC TO

NKDA

Date: _____ **Estimated Length of Stay:** _____

ORDER PATIENT STATUS: **Observation Indication** _____ (e.g., nausea, bleeding) **Outpatient in a bed**
(additional time needed to evaluate patient's condition) (Normal recovery requiring acute care bed)

Inpatient (meets medical necessity for admission) **Outpatient procedure**

Hospitalization certified necessary for the following reasons:

Plan of Care: See Progress Notes, H and P, and Physician Orders

D/C Plan: Home with Office Follow-up Home Health Care Rehabilitation Care Facility
 Extended Care Facility (e.g. Adult Home, Nursing Home, etc.) Other (Specify): _____

Physician Signature: _____

ORDERS (other than medications) CHECK ALL THAT APPLY	MEDICATION & I.V. FLUID ORDERS CHECK ALL THAT APPLY
<ol style="list-style-type: none"> 1. Patient must be in critical care or step down unit 2. Baseline PTT, PT/INR, Creatinine, CMP, HCT, Platelet Count 3. Platelet count every day 4. All PTTs are to be run stat 5. PTT 2 hr and 6 hr after initiation of Argatroban 6. PTT 4 hr after any dosage change until two consecutive PTTs are therapeutic (40-83 seconds) then change PTT to q day 7. If PTT is greater than 100 seconds stop argatroban, draw PTT every 2 hours until PTT is 40-83 seconds then restart Argatroban at new rate (see Rate Chart) 8. Daily PT/INR after warfarin started as noted below <ol style="list-style-type: none"> a. If Argatroban dose is greater than 2 mcg/kg/min, reduce Argatroban to 2 mcg/kg/min for 4 hours, draw PT/INR. If PT/INR greater than 4, on Warfarin and Argatroban, stop Argatroban, draw PT/INR four hours later, restart Argatroban immediately after PT/INR drawn at original rate. Use second PT/INR for Warfarin dosage adjustments. b. If Argatroban dose is less than or equal to 2 mcg/kg/min drawn PT/INR, if PT/INR greater than 4, on Warfarin and Argatroban, stop Argatroban, draw PT/INR four hours later, restart Argatroban immediately after PT/INR drawn at prior rate. Use second PT/INR for Warfarin dosage adjustments. 9. Guaiac all stools, gastrocuccult all emesis, visually check for hematuria or other bleeding. 10. Notify physician immediately for unexplained drop in blood pressure, development of hematoma, drop in HCT, flank pain, bright red urine, bruising or other signs of bleeding. 	<p>Discontinue all Heparin, Heparin flushes, Enoxaparin (Lovenox) and Dalteparin (Fragmin)</p> <p>Discontinue Warfarin (Coumadin)</p> <p>Phytonadione (Vitamin K) 10 mg subQ x 1 (if the patient has been on Warfarin)</p> <p>Discontinue all IM injections (physician to give order for alternate route of administration)</p> <p>Argatroban infusion:</p> <p><input type="checkbox"/> 2 mcg/kg/min if hepatic disease score per table is less than or equal to 6</p> <p><input type="checkbox"/> 1 mcg/kg/min if creatinine clearance is less than 60 ml/min and hepatic disease score is less than or equal to 6.</p> <p><input type="checkbox"/> 0.5 mcg/kg/min if hepatic disease score per table is greater than 6</p> <p>Adjust Argatroban rate per PTT using Argatroban rate adjustment chart</p> <p>Pharmacy to send initial infusion rate chart and rate adjustment chart to floor with Argatroban infusion</p> <p>Warfarin (Coumadin) _____ mg (range 2 - 5 mg) Qday once platelet count is greater than 150,000 per mcL Goal PT/INR while argatroban infusing at or below 2 mcg/kg/min is 4-5, off argatroban goal PT/INR is 2-3.</p> <p>Discontinue Argatroban when two consecutive daily INR are 2 to 3 (on Warfarin alone, see step 8 on left) and Warfarin therapy has overlapped Argatroban therapy for at least 5 days.</p>

DO NOT USE ABBREVIATIONS:			
MS	U (unit)	IU (International Unit)	Q.D. QD q.d. qd (daily)
MISO4	Trailing zero (X.0mg)		Q.O.D. QOD q.o.d. qod (every other day)
MgSO4	Lack of leading zero (.Xmg)		

Physician Signature _____

Date / Time _____



BON SECOURS
RICHMOND HEALTH SYSTEM

- Memorial Regional Medical Center
- Richmond Community Hospital
- St. Francis Medical Center
- St. Mary's Hospital

Place patient label inside box (if no patient label, complete below)

Name: _____

DOB: _____

MR #: _____

**ARGATROBAN PROTOCOL
FOR HIT/ISOLATED HIT**

ALL ORDERS MUST INCLUDE DATE, TIME,
AND PHYSICIAN'S SIGNATURE.

THIS PATIENT IS ALLERGIC TO

NKDA

Date: _____ **Estimated Length of Stay:** _____

ORDER PATIENT STATUS: **Observation Indication** _____ (e.g., nausea, bleeding) **Outpatient in a bed**
(additional time needed to evaluate patient's condition) (Normal recovery requiring acute care bed)

Inpatient (meets medical necessity for admission) **Outpatient procedure**

Hospitalization certified necessary for the following reasons:

Plan of Care: See Progress Notes, H and P, and Physician Orders

D/C Plan: Home with Office Follow-up Home Health Care Rehabilitation Care Facility
 Extended Care Facility (e.g. Adult Home, Nursing Home, etc.) Other (Specify): _____

Physician Signature: _____

ORDERS (other than medications) CHECK ALL THAT APPLY	MEDICATION & I.V. FLUID ORDERS CHECK ALL THAT APPLY
<ol style="list-style-type: none"> 1. Patient must be in critical care or step down unit 2. Baseline PTT, PT/INR, Creatinine, CMP, HCT, Platelet Count 3. Platelet count every day 4. All PTTs are to be run stat 5. PTT 2 hr and 6 hr after initiation of Argatroban 6. PTT 4 hr after any dosage change until two consecutive PTTs are therapeutic (40-83 seconds) then change PTT to q day 7. If PTT is greater than 100 seconds stop argatroban, draw PTT every 2 hours until PTT is 40-83 seconds then restart Argatroban at new rate (see Rate Chart) 8. Daily PT/INR after warfarin started as noted below <ol style="list-style-type: none"> a. If Argatroban dose is greater than 2 mcg/kg/min, reduce Argatroban to 2 mcg/kg/min for 4 hours, draw PT/INR. If PT/INR greater than 4, on Warfarin and Argatroban, stop Argatroban, draw PT/INR four hours later, restart Argatroban immediately after PT/INR drawn at original rate. Use second PT/INR for Warfarin dosage adjustments. b. If Argatroban dose is less than or equal to 2 mcg/kg/min drawn PT/INR, if PT/INR greater than 4, on Warfarin and Argatroban, stop Argatroban, draw PT/INR four hours later, restart Argatroban immediately after PT/INR drawn at prior rate. Use second PT/INR for Warfarin dosage adjustments. 9. Guaiac all stools, gastrocuccult all emesis, visually check for hematuria or other bleeding. 10. Notify physician immediately for unexplained drop in blood pressure, development of hematoma, drop in HCT, flank pain, bright red urine, bruising or other signs of bleeding. 	<p>Discontinue all Heparin, Heparin flushes, Enoxaparin (Lovenox) and Dalteparin (Fragmin)</p> <p>Discontinue Warfarin (Coumadin)</p> <p>Phytonadione (Vitamin K) 10 mg subQ x 1 (if the patient has been on Warfarin)</p> <p>Discontinue all IM injections (physician to give order for alternate route of administration)</p> <p>Argatroban infusion:</p> <p><input type="checkbox"/> 2 mcg/kg/min if hepatic disease score per table is less than or equal to 6</p> <p><input type="checkbox"/> 1 mcg/kg/min if creatinine clearance is less than 60 ml/min and hepatic disease score is less than or equal to 6.</p> <p><input type="checkbox"/> 0.5 mcg/kg/min if hepatic disease score per table is greater than 6</p> <p>Adjust Argatroban rate per PTT using Argatroban rate adjustment chart</p> <p>Pharmacy to send initial infusion rate chart and rate adjustment chart to floor with Argatroban infusion</p> <p>Warfarin (Coumadin) _____ mg (range 2 - 5 mg) Qday once platelet count is greater than 150,000 per mcL Goal PT/INR while argatroban infusing at or below 2 mcg/kg/min is 4-5, off argatroban goal PT/INR is 2-3.</p> <p>Discontinue Argatroban when two consecutive daily INR are 2 to 3 (on Warfarin alone, see step 8 on left) and Warfarin therapy has overlapped Argatroban therapy for at least 5 days.</p>

DO NOT USE ABBREVIATIONS:			
MS	U (unit)	IU (International Unit)	Q.D. QD q.d. qd (daily)
MISO4	Trailing zero (X.0mg)		Q.O.D. QOD q.o.d. qod (every other day)
MgSO4	Lack of leading zero (.Xmg)		

Physician Signature

Date / Time



BON SECOURS
RICHMOND HEALTH SYSTEM

- Memorial Regional Medical Center
- Richmond Community Hospital
- St. Francis Medical Center
- St. Mary's Hospital

Place patient label inside box (if no patient label, complete below)

Name: _____

DOB: _____

MR #: _____

**ARGATROBAN PROTOCOL
FOR HIT/ISOLATED HIT**

ALL ORDERS MUST INCLUDE DATE, TIME,
AND PHYSICIAN'S SIGNATURE.

THIS PATIENT IS ALLERGIC TO

NKDA

ORDERS (other than medications) CHECK ALL THAT APPLY	MEDICATION & I.V. FLUID ORDERS CHECK ALL THAT APPLY																																
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th colspan="4" style="text-align: center;">Hepatic Disease Score for Argatroban Dosing</th> </tr> <tr> <th></th> <th style="text-align: center;">1 point</th> <th style="text-align: center;">2 points</th> <th style="text-align: center;">3 points</th> </tr> </thead> <tbody> <tr> <td>Bilirubin (mg/dl)</td> <td style="text-align: center;">less than 2</td> <td style="text-align: center;">2-3</td> <td style="text-align: center;">greater than 3</td> </tr> <tr> <td></td> <td style="text-align: center;">less than 4 for patients with primary biliary cirrhosis</td> <td style="text-align: center;">4-10 for patients with primary biliary cirrhosis</td> <td style="text-align: center;">greater than 10 for patients with primary biliary cirrhosis</td> </tr> <tr> <td>Albumin (g/dl)</td> <td style="text-align: center;">greater than 3.5</td> <td style="text-align: center;">2.8-3.5</td> <td style="text-align: center;">less than 2.8</td> </tr> <tr> <td>PT prolongation (INR)</td> <td style="text-align: center;">less than 4 seconds (INR less than 1.7)</td> <td style="text-align: center;">4-6 seconds (INR 1.7-2.3)</td> <td style="text-align: center;">greater than 6 seconds (INR greater than 2.3)</td> </tr> <tr> <td>Ascites</td> <td style="text-align: center;">Absent</td> <td style="text-align: center;">Slight</td> <td style="text-align: center;">Moderate</td> </tr> <tr> <td>Encephalopat by</td> <td style="text-align: center;">Absent</td> <td style="text-align: center;">Mild (1-2)</td> <td style="text-align: center;">Severe (3-4)</td> </tr> </tbody> </table> <p>To calculate the Hepatic Disease Score, determine the individual score (1,2,3) for each parameter. If the sum of all parameters is greater than 6 then the patient should be started on 0.5 mcg/kg/min of Argatroban.</p> <p>Stages of Hepatic Encephalopathy</p> <p>Stage 1: Euphoria or depression, mild confusion, slurred speech, disordered sleep</p> <p>Stage 2: Lethargy, moderate confusion</p> <p>Stage 3: Marked confusion, incoherent speech, sleeping but arousable</p> <p>Stage 4: Coma, initially responsive to noxious stimuli, but later unresponsive</p>	Hepatic Disease Score for Argatroban Dosing					1 point	2 points	3 points	Bilirubin (mg/dl)	less than 2	2-3	greater than 3		less than 4 for patients with primary biliary cirrhosis	4-10 for patients with primary biliary cirrhosis	greater than 10 for patients with primary biliary cirrhosis	Albumin (g/dl)	greater than 3.5	2.8-3.5	less than 2.8	PT prolongation (INR)	less than 4 seconds (INR less than 1.7)	4-6 seconds (INR 1.7-2.3)	greater than 6 seconds (INR greater than 2.3)	Ascites	Absent	Slight	Moderate	Encephalopat by	Absent	Mild (1-2)	Severe (3-4)	
Hepatic Disease Score for Argatroban Dosing																																	
	1 point	2 points	3 points																														
Bilirubin (mg/dl)	less than 2	2-3	greater than 3																														
	less than 4 for patients with primary biliary cirrhosis	4-10 for patients with primary biliary cirrhosis	greater than 10 for patients with primary biliary cirrhosis																														
Albumin (g/dl)	greater than 3.5	2.8-3.5	less than 2.8																														
PT prolongation (INR)	less than 4 seconds (INR less than 1.7)	4-6 seconds (INR 1.7-2.3)	greater than 6 seconds (INR greater than 2.3)																														
Ascites	Absent	Slight	Moderate																														
Encephalopat by	Absent	Mild (1-2)	Severe (3-4)																														

DO NOT USE ABBREVIATIONS:
 MS U (unit) IU (International Unit) Q.D. QD q.d. qd (daily)
 MSO4 Trailing zero (X.0mg) Q.O.D. QOD q.o.d. qod (every other day)
 MgSO4 Lack of leading zero (.Xmg)

Physician Signature

Date / Time



BON SECOURS
RICHMOND HEALTH SYSTEM

- Memorial Regional Medical Center
- Richmond Community Hospital
- St. Francis Medical Center
- St. Mary's Hospital

Place patient label inside box (if no patient label, complete below)

Name: _____
 DOB: _____
 MR #: _____

**ARGATROBAN PROTOCOL
FOR HIT/ISOLATED HIT**

ALL ORDERS MUST INCLUDE DATE, TIME,
AND PHYSICIAN'S SIGNATURE.

THIS PATIENT IS ALLERGIC TO

NKDA

ORDERS (other than medications) CHECK ALL THAT APPLY	MEDICATION & I.V. FLUID ORDERS CHECK ALL THAT APPLY																																
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th colspan="4" style="text-align: center;">Hepatic Disease Score for Argatroban Dosing</th> </tr> <tr> <th></th> <th style="text-align: center;">1 point</th> <th style="text-align: center;">2 points</th> <th style="text-align: center;">3 points</th> </tr> </thead> <tbody> <tr> <td>Bilirubin (mg/dl)</td> <td style="text-align: center;">less than 2</td> <td style="text-align: center;">2-3</td> <td style="text-align: center;">greater than 3</td> </tr> <tr> <td></td> <td style="text-align: center;">less than 4 for patients with primary biliary cirrhosis</td> <td style="text-align: center;">4-10 for patients with primary biliary cirrhosis</td> <td style="text-align: center;">greater than 10 for patients with primary biliary cirrhosis</td> </tr> <tr> <td>Albumin (g/dl)</td> <td style="text-align: center;">greater than 3.5</td> <td style="text-align: center;">2.8-3.5</td> <td style="text-align: center;">less than 2.8</td> </tr> <tr> <td>PT prolongation (INR)</td> <td style="text-align: center;">less than 4 seconds (INR less than 1.7)</td> <td style="text-align: center;">4-6 seconds (INR 1.7-2.3)</td> <td style="text-align: center;">greater than 6 seconds (INR greater than 2.3)</td> </tr> <tr> <td>Ascites</td> <td style="text-align: center;">Absent</td> <td style="text-align: center;">Slight</td> <td style="text-align: center;">Moderate</td> </tr> <tr> <td>Encephalopat by</td> <td style="text-align: center;">Absent</td> <td style="text-align: center;">Mild (1-2)</td> <td style="text-align: center;">Severe (3-4)</td> </tr> </tbody> </table> <p>To calculate the Hepatic Disease Score, determine the individual score (1,2,3) for each parameter. If the sum of all parameters is greater than 6 then the patient should be started on 0.5 mcg/kg/min of Argatroban.</p> <p>Stages of Hepatic Encephalopathy</p> <p>Stage 1: Euphoria or depression, mild confusion, slurred speech, disordered sleep</p> <p>Stage 2: Lethargy, moderate confusion</p> <p>Stage 3: Marked confusion, incoherent speech, sleeping but arousable</p> <p>Stage 4: Coma, initially responsive to noxious stimuli, but later unresponsive</p>	Hepatic Disease Score for Argatroban Dosing					1 point	2 points	3 points	Bilirubin (mg/dl)	less than 2	2-3	greater than 3		less than 4 for patients with primary biliary cirrhosis	4-10 for patients with primary biliary cirrhosis	greater than 10 for patients with primary biliary cirrhosis	Albumin (g/dl)	greater than 3.5	2.8-3.5	less than 2.8	PT prolongation (INR)	less than 4 seconds (INR less than 1.7)	4-6 seconds (INR 1.7-2.3)	greater than 6 seconds (INR greater than 2.3)	Ascites	Absent	Slight	Moderate	Encephalopat by	Absent	Mild (1-2)	Severe (3-4)	
Hepatic Disease Score for Argatroban Dosing																																	
	1 point	2 points	3 points																														
Bilirubin (mg/dl)	less than 2	2-3	greater than 3																														
	less than 4 for patients with primary biliary cirrhosis	4-10 for patients with primary biliary cirrhosis	greater than 10 for patients with primary biliary cirrhosis																														
Albumin (g/dl)	greater than 3.5	2.8-3.5	less than 2.8																														
PT prolongation (INR)	less than 4 seconds (INR less than 1.7)	4-6 seconds (INR 1.7-2.3)	greater than 6 seconds (INR greater than 2.3)																														
Ascites	Absent	Slight	Moderate																														
Encephalopat by	Absent	Mild (1-2)	Severe (3-4)																														

DO NOT USE ABBREVIATIONS:
 MS U (unit) IU (International Unit) Q.D. QD q.d. qd (daily)
 MSO4 Trailing zero (X.0mg) Q.O.D. QOD q.o.d. qod (every other day)
 MgSO4 Lack of leading zero (.Xmg)

Physician Signature

Date / Time



BON SECOURS
RICHMOND HEALTH SYSTEM

- Memorial Regional Medical Center
- Richmond Community Hospital
- St. Francis Medical Center
- St. Mary's Hospital

Place patient label inside box (if no patient label, complete below)

Name: _____

DOB: _____

MR #: _____

**ARGATROBAN PROTOCOL
FOR HIT/ISOLATED HIT**

ALL ORDERS MUST INCLUDE DATE, TIME,
AND PHYSICIAN'S SIGNATURE.

THIS PATIENT IS ALLERGIC TO

NKDA

ORDERS (other than medications) CHECK ALL THAT APPLY	MEDICATION & I.V. FLUID ORDERS CHECK ALL THAT APPLY																																
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th colspan="4" style="text-align: center;">Hepatic Disease Score for Argatroban Dosing</th> </tr> <tr> <th></th> <th style="text-align: center;">1 point</th> <th style="text-align: center;">2 points</th> <th style="text-align: center;">3 points</th> </tr> </thead> <tbody> <tr> <td>Bilirubin (mg/dl)</td> <td style="text-align: center;">less than 2</td> <td style="text-align: center;">2-3</td> <td style="text-align: center;">greater than 3</td> </tr> <tr> <td></td> <td style="text-align: center;">less than 4 for patients with primary biliary cirrhosis</td> <td style="text-align: center;">4-10 for patients with primary biliary cirrhosis</td> <td style="text-align: center;">greater than 10 for patients with primary biliary cirrhosis</td> </tr> <tr> <td>Albumin (g/dl)</td> <td style="text-align: center;">greater than 3.5</td> <td style="text-align: center;">2.8-3.5</td> <td style="text-align: center;">less than 2.8</td> </tr> <tr> <td>PT prolongation (INR)</td> <td style="text-align: center;">less than 4 seconds (INR less than 1.7)</td> <td style="text-align: center;">4-6 seconds (INR 1.7-2.3)</td> <td style="text-align: center;">greater than 6 seconds (INR greater than 2.3)</td> </tr> <tr> <td>Ascites</td> <td style="text-align: center;">Absent</td> <td style="text-align: center;">Slight</td> <td style="text-align: center;">Moderate</td> </tr> <tr> <td>Encephalopat by</td> <td style="text-align: center;">Absent</td> <td style="text-align: center;">Mild (1-2)</td> <td style="text-align: center;">Severe (3-4)</td> </tr> </tbody> </table> <p>To calculate the Hepatic Disease Score, determine the individual score (1,2,3) for each parameter. If the sum of all parameters is greater than 6 then the patient should be started on 0.5 mcg/kg/min of Argatroban.</p> <p>Stages of Hepatic Encephalopathy</p> <p>Stage 1: Euphoria or depression, mild confusion, slurred speech, disordered sleep</p> <p>Stage 2: Lethargy, moderate confusion</p> <p>Stage 3: Marked confusion, incoherent speech, sleeping but arousable</p> <p>Stage 4: Coma, initially responsive to noxious stimuli, but later unresponsive</p>	Hepatic Disease Score for Argatroban Dosing					1 point	2 points	3 points	Bilirubin (mg/dl)	less than 2	2-3	greater than 3		less than 4 for patients with primary biliary cirrhosis	4-10 for patients with primary biliary cirrhosis	greater than 10 for patients with primary biliary cirrhosis	Albumin (g/dl)	greater than 3.5	2.8-3.5	less than 2.8	PT prolongation (INR)	less than 4 seconds (INR less than 1.7)	4-6 seconds (INR 1.7-2.3)	greater than 6 seconds (INR greater than 2.3)	Ascites	Absent	Slight	Moderate	Encephalopat by	Absent	Mild (1-2)	Severe (3-4)	
Hepatic Disease Score for Argatroban Dosing																																	
	1 point	2 points	3 points																														
Bilirubin (mg/dl)	less than 2	2-3	greater than 3																														
	less than 4 for patients with primary biliary cirrhosis	4-10 for patients with primary biliary cirrhosis	greater than 10 for patients with primary biliary cirrhosis																														
Albumin (g/dl)	greater than 3.5	2.8-3.5	less than 2.8																														
PT prolongation (INR)	less than 4 seconds (INR less than 1.7)	4-6 seconds (INR 1.7-2.3)	greater than 6 seconds (INR greater than 2.3)																														
Ascites	Absent	Slight	Moderate																														
Encephalopat by	Absent	Mild (1-2)	Severe (3-4)																														

DO NOT USE ABBREVIATIONS:
 MS U (unit) IU (International Unit) Q.D. QD q.d. qd (daily)
 MSO4 Trailing zero (X.0mg) Q.O.D. QOD q.o.d. qod (every other day)
 MgSO4 Lack of leading zero (.Xmg)

Physician Signature

Date / Time